



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Now we have

$$\begin{aligned}
 6^\lambda \Sigma N(d) N(m/d) &= 6^\lambda \sum_{r_1=0}^{\alpha_1} (\alpha_1 - r_1 + 1)(\alpha_2 - r_2 + 1) \cdots (\alpha_\lambda - r_\lambda + 1)(r_1 + 1)(r_2 + 1) \\
 &\quad \cdots (r_\lambda + 1) \\
 &= 6^\lambda \sum_{r_1=0}^{\alpha_1} (\alpha_1 - r_1 + 1)(r_1 + 1) \cdots (\alpha_\lambda - r_\lambda + 1)(r_\lambda + 1) \\
 &= 6 \sum_{r_1=0}^{\alpha_1} [\alpha_1(r_1 + 1) - (r_1^2 - 1)] \cdot 6 \sum_{r_2=0}^{\alpha_2} [\alpha_2(r_2 + 1) - (r_2^2 - 1)] \\
 &\quad \cdots 6 \sum_{r_\lambda=0}^{\alpha_\lambda} [\alpha_\lambda(r_\lambda + 1) - (r_\lambda^2 - 1)].
 \end{aligned} \tag{2}$$

The indicated summations can be readily performed. We have

$$\sum_{r_i=0}^{\alpha_i} (r_i + 1) = \frac{\alpha_i(\alpha_i + 1)}{2} + \alpha_i + 1 = \frac{(\alpha_i + 1)(\alpha_i + 2)}{2}$$

and

$$\sum_{r_i=0}^{\alpha_i} (r_i^2 - 1) = \frac{\alpha_i(\alpha_i + 1)(2\alpha_i + 1)}{6} - \alpha_i - 1 = \frac{(\alpha_i + 1)(\alpha_i + 2)(2\alpha_i - 3)}{6}.$$

Hence

$$\begin{aligned}
 6 \sum_{r_i=0}^{\alpha_i} [\alpha_i(r_i + 1) - (r_i^2 - 1)] &= 6 \left[\alpha_i \frac{(\alpha_i + 1)(\alpha_i + 2)}{2} - \frac{(\alpha_i + 1)(\alpha_i + 2)(2\alpha_i - 3)}{6} \right] \\
 &= (\alpha_i + 1)(\alpha_i + 2)(\alpha_i + 3).
 \end{aligned}$$

Substituting for these sums in the last member of (2) and making use of (1), we have

$$\begin{aligned}
 6^\lambda \Sigma N(d) N(m/d) &= (\alpha_1 + 1)(\alpha_1 + 2)(\alpha_1 + 3)(\alpha_2 + 1)(\alpha_2 + 2)(\alpha_2 + 3) \\
 &\quad \cdots (\alpha_\lambda + 1)(\alpha_\lambda + 2)(\alpha_\lambda + 3) \\
 &= N(m) N(Pm) N(P^2m).
 \end{aligned}$$

197. Proposed by E. T. BELL, Seattle, Washington.

Show that in the expansion of

$$\frac{1 + z + z^2 + \cdots + z^{p-1}}{(1 - z)^{p-1}} - 1,$$

where p is a prime, the coefficients of the various powers of z are divisible by p .
[Eisenstein, *Crelle*, t. 27, p. 282.]

SOLUTION BY B. F. YANNEY, University of Wooster.

The given expression is evidently equal to $(1 - z^p)/(1 - z)^p - 1$, which may be put in the form $(1 - z^p)(1 - z)^{-p} - 1$. Expanding the second factor of the first term and noticing that, since p is prime, the coefficient of each term

in the expansion except that of the 1st, the $(p+1)$ th, the $(2p+1)$ th, and so on to the $(np+1)$ th, and so on, is divisible by p , we obtain, omitting the terms with the divisible coefficients,

$$(1 - z^p) \left(1 + \frac{p(p+1) \cdots (2p-1)}{p!} z^p + \frac{p(p+1) \cdots (3p-1)}{(2p)!} z^{2p} + \cdots \right. \\ \left. + \frac{p(p+1) \cdots ((n+1)p-1)}{(np)!} \cdots \right) - 1.$$

Performing the indicated operations, we get

$$\left(\frac{p(p+1) \cdots (2p-1)}{p!} - 1 \right) z^p + \left(\frac{p(p+1) \cdots (3p-1)}{(2p)!} - \frac{p(p+1) \cdots (2p-1)}{p!} \right) z^{2p} \\ + \cdots + \left(\frac{p(p+1) \cdots ((n+1)p-1)}{(np)!} - \frac{p(p+1) \cdots (np-1)}{((n-1)p)!} \right) z^{np} + \cdots.$$

The first coefficient may be written $[(p+1)(p+2) \cdots (p+p-1) - (p-1)!]/(p-1)!$, which is equal to $[pA + (p-1)! - (p-1)!]/(p-1)!$, where A is a polynomial in p . This expression, which is equal to $pA/(p-1)!$, and is an integer, as are all the coefficients, is plainly divisible by p , since the denominator does not contain p as a factor.

The coefficient of the general term may be changed in form to

$$\frac{p(p+1) \cdots (np-1)}{((n-1)p)!} \left(\frac{np(np+1) \cdots (np+p-1)}{[(n-1)p+1][(n-1)p+2] \cdots [np]} - 1 \right).$$

The first factor of this is an integer; hence so is the second, which latter can be expressed in the form

$$\frac{(np+1)(np+2) \cdots (np+p-1) - [(n-1)p+1][(n-1)p+2] \cdots [(n-1)p+p-1]}{[(n-1)p+1][(n-1)p+2] \cdots [(n-1)p+p-1]},$$

which equals

$$\frac{(npB + (p-1)!) - ((n-1)pC + (p-1)!)}{(n-1)pC + (p-1)!},$$

where B and C are polynomials in np and $(n-1)p$, respectively. Since the denominator does not contain p as a factor, but the numerator does, the expression is divisible by p , which completes the proof.

Also solved by ELMER SCHUYLER, H. C. FEEMSTER and the PROPOSER.